arXiv:1301.2828v2 [math.PR] 24 Jan 2013

On the nonuniform Berry–Esseen bound

Iosif Pinelis[*]

Department of Mathematical Sciences
Michigan Technological University
Houghton, Michigan 49931, USA
E-mail: `ipinelis@mtu.edu`

Abstract: Due to the effort of a number of authors, the value c_{u} of the absolute constant factor in the uniform Berry–Esseen (BE) bound for sums of independent random variables has been gradually reduced to 0.4748 in the iid case and 0.5600 in the general case; both these values were recently obtained by Shevtsova. On the other hand, Esseen had shown that c_{u} cannot be less than 0.4097. Thus, the gap factor between the best known upper and lower bounds on (the least possible value of) c_{u} is now rather close to 1.

The situation is quite different for the absolute constant factor c_{nu} in the corresponding nonuniform BE bound. Namely, the best correctly established upper bound on c_{nu} in the iid case is about 30 times the corresponding best known lower bound, and this gap factor is greater than 100 in the general case. In the present paper, improvements to the prevailing method (going back to S. Nagaev) of obtaining nonuniform BE bounds are suggested. Moreover, a new method is presented, of a rather purely Fourier kind, based on a family of smoothing inequalities, which work better in the tail zones. As an illustration, a quick proof of Nagaev's nonuniform BE bound is given. Some further refinements in the application of the method are shown as well.

AMS 2010 subject classifications: 60E15, 62E17.
Keywords and phrases: Berry–Esseen bounds, rate of convergence to normality, probability inequalities, smoothing inequalities, sums of independent random variables.

Contents

1 Uniform and nonuniform Berry–Esseen (BE) bounds . . . 1
2 The Bohman–Prawitz–Vaaler smoothing inequalities . . . 2
3 Nonuniform BE bounds: Nagaev's result and method . . . 3
  3.1 A historical sketch of the problem of nonuniform BE bounds . . . 5
  3.2 Possible improvements of Nagaev's method . . . 6
4 A new way to obtain nonuniform BE bounds . . . 9
5 Constructions of the smoothing filter M . . . 14
References . . . 18

1. Uniform and nonuniform Berry–Esseen (BE) bounds

Suppose that $X_1, \dots, X_n$ are independent zero-mean r.v.'s, with

$$S := X_1 + \dots + X_n,\ A := \sum \mathsf{E}\,|X_i|^3 < \infty,\ \text{and } B := \sqrt{\sum \mathsf{E}\,|X_i|^2} > 0.$$

Consider

$$\Delta(z) := |\,\mathsf{P}(S > Bz) - \mathsf{P}(Z > z)| \quad \text{and} \quad r_L := A/B^3,$$

where $Z \sim N(0,1)$ and $z \geqslant 0$; of course, r_L is the so-called Lyapunov ratio. Note that, in the "iid" case (when the X_i's are iid), r_L will be on the order of $1/\sqrt{n}$.

---

[*] Supported by NSA grant H98230-12-1-0237

In such an iid case, let us also assume that $\mathsf{E}\,X_1^2 = 1$.

Uniform and nonuniform BE bounds are upper bounds on $\Delta(z)$ of the forms

$$c_{\mathsf{u}}\, r_L \quad \text{and} \quad c_{\mathsf{nu}} \frac{r_L}{1+z^3}, \tag{1.1}$$

respectively, for some absolute positive real constants c_{u} and c_{nu} and for all $z \geqslant 0$.

Apparently the best currently known upper bound on c_{u} (in the iid case) is due to Shevtsova [39] and is given by the inequality

$$c_{\mathsf{u}} \leqslant 0.4748. \tag{1.2}$$

On the other hand, Esseen's example [8] with iid X_i's, $n \to \infty$, z appropriately close to 0, and

$$\mathsf{P}(X_1 = 1 - p_{\mathsf{Ess}}) = p_{\mathsf{Ess}} = 1 - \mathsf{P}(X_1 = -p_{\mathsf{Ess}}) \tag{1.3}$$

with $p_{\mathsf{Ess}} := 2 - \sqrt{10}/2 = 0.4188...$ showed that c_{u} cannot be less than $\frac{3+\sqrt{10}}{6\sqrt{2\pi}} = 0.4097\dots$; a similar lower bound on the BE constant for intervals was recently shown by Dinev and Mattner [6] to be $\sqrt{\frac{2}{\pi}} = 0.7978\dots$, which is almost twice as large as $0.4097\dots$.

Thus, the optimal value of c_{u} is already known to be within the rather small interval from 0.4097 to 0.4748 in the iid case $\big($in the general case the best known upper bound on c_{u} appears to be 0.5600, due to Shevtsova [40]; a slightly worse upper bound, 0.5606, is due to Tyurin [44]$\big)$.

2. The Bohman–Prawitz–Vaaler smoothing inequalities

To a significant extent the mentioned best known uniform BE bounds are based on the smoothing result due to Prawitz [38, (1a, 1b)], which states the following. There exists a nonempty class of functions $M\colon \mathbb{R} \to \mathbb{C}$ such that

$$M(t) = 0 \quad \text{if} \quad |t| > 1 \tag{2.1}$$

and for any r.v. X, any real $T > 0$, and any real x,

$$\mathfrak{G}\big(M_T(-\#)\,\mathsf{E}\,e^{iX\#}\big)(x) \leqslant \mathsf{P}(X < x) - \tfrac12 \leqslant \mathsf{P}(X \leqslant x) - \tfrac12 \leqslant \mathfrak{G}\big(M_T(\#)\,\mathsf{E}\,e^{iX\#}\big)(x), \tag{2.2}$$

where

$$M_T(\#) := M(\#/T), \tag{2.3}$$

$$\mathfrak{G}(f)(x) := \frac{i}{2\pi}\ \text{p.v.} \int_{\infty}^{\infty} e^{-itx} f(t) \frac{\mathrm{d}t}{t}, \tag{2.4}$$

and p.v. stands for "principal value", so that p.v. $\int_{-\infty}^{\infty} := \lim_{\substack{\varepsilon\downarrow 0\\ A\uparrow\infty}} \big(\int_{-A}^{-\varepsilon} + \int_{\varepsilon}^{A}\big)$; here and subsequently, the symbol # stands for the argument of a function. Of course, the upper and lower bounds in (2.2) must take on only real values; this can be provided by the condition that

$$M_1 := \operatorname{Re} M \text{ is even} \quad \text{and} \quad M_2 := \operatorname{Im} M \text{ is odd}. \tag{2.5}$$

Note also that the upper and lower bounds in (2.2) easily follow from each other, by changing X to $-X$.

Inequalities (2.2) may be compared with the corresponding well-known inversion formula

$$\mathsf{P}(X < x) + \tfrac12\,\mathsf{P}(X = x) - \tfrac12 = \mathfrak{G}(\mathsf{E}\,e^{iX\#})(x) \tag{2.6}$$

for all real x; see e.g. [10, (2)]. The multiplier $M(\#)$ of the c.f. $\mathsf{E}\, e^{iX\#}$ in (2.2) is the Fourier transform of the function $\check{M}(\#) := \frac{1}{2\pi}\int_{-\infty}^{\infty} e^{-it\#} M(t)\,\mathrm{d}t$, which may be considered as a smoothing kernel – since, in view of (2.1), the spectral decomposition of $\check{M}$ does not have components of frequencies greater than 1. So, the factors $M(\pm\#/T)$ in the bounds in (2.2) *filter* out the components of the function $\mathfrak{G}(\mathsf{E}\, e^{iX\#})$ of frequencies greater than T and thus make the function smoother and flatter, especially if T is not large enough. Another way to look at such smoothing is through the Paley–Wiener theory, which implies that the Fourier spectrum of a function is contained in the interval $[-T,T]$ iff the function is (the restriction to $\mathbb{R}$ of) an entire analytic function of exponential type T and hence rather slowly varying if T is not large; see e.g. [7, Section 43]. On the other hand, from an analytical viewpoint, the presence of the factors $M(\pm\#/T)$ is useful, because one then needs to bound the values $\mathsf{E}\, e^{itX}$ of the c.f. of X only for $t \in [-T,T]$, which is a much easier task unless T is too large.

One particular function M for which (2.2) holds is given by the formula

$$M(t) = \big[(1-|t|)\,\pi t\cot\pi t + |t| - i(1-|t|)\,\pi t\big]\,\mathrm{I}\{|t|<1\} \tag{2.7}$$

for all $t \neq 0$ [38]; here and subsequently, it is tacitly assumed that the functions of interest are extended to 0 by continuity. For this particular multiplier M, which was shown in [38] to have a certain optimality property, the corresponding smoothing kernel $\check{M}(\#) := \frac{1}{2\pi}\int_{\mathbb{R}} e^{-it\#} M(t)\,\mathrm{d}t$ is given by the formula

$$\check{M}(x) = \frac{2\pi x\sin x\,\big(2\pi(x+2\pi) - x^2\psi'\big(\frac{x}{2\pi}\big)\big) - (1-\cos x)\,\big(x^3\psi''\big(\frac{x}{2\pi}\big) + 4\pi^2(x+4\pi)\big)}{4\pi^3x^3}$$

for $x \neq 0$, where ψ is the digamma function, defined by the formula $\psi(z) = \Gamma'(z)/\Gamma(z)$; this kernel is (necessarily) asymmetric and alternating in sign; also, $\int_{-\infty}^{\infty}\check{M}(x)\,\mathrm{d}x = M(0) = 1$; a part of the graph of this kernel $\check{M}$ is shown here on the left.



Earlier, inequalities of the form (2.2) were obtained by Bohman [3] for another class of functions M, with apparently not quite as good approximation properties. Another approach to Prawitz's results was demonstrated by Vaaler [45].

3. Nonuniform BE bounds: Nagaev's result and method

The classical result by Nagaev [19] is that in the "iid" case

$$|\,\mathsf{P}(S > z\sqrt{n}) - \mathsf{P}(Z > z)| \leqslant c_{\mathsf{nu}}\frac{\mathsf{E}\,|X_1|^3}{(1+z^3)\sqrt{n}} \tag{3.1}$$

for all real $z \geqslant 0$, where c_{nu} is an absolute constant. Bikelis [2] extended this result to the case of non-iid X_i's. Nagaev's method involves the following essential components:

- truncation;
- Cramer's exponential tilt, together with a uniform BE bound;
- an exponential bound on large deviation probabilities.

First, truncated versions of X_i, say $X_i^{(y)}$, are obtained, such that $X_i^{(y)} \leqslant y$ for some real $y > 0$ and all i $\big($the r.v.'s $X_i^{(y)}$ may, in some variants of this approach including [19], be improper in the sense that they may take values that are not real numbers$\big)$. The truncation is done in order to make the exponential tilt and an exponential inequality possible. The value of the truncation level y is chosen (i) to be large enough so that the tails of the truncated sum $S^{(y)} := X_1^{(y)} + \cdots + X_n^{(y)}$ be close enough to those of S and, on the other hand, (ii) to be small enough so that the exponential tilt and the exponential inequality result in not too large a bound. In some variants, including the ones in [19, 2], two different truncation levels are used.

In view of the uniform BE bound, without loss of generality $z \geqslant z_0$, where z_0 is an arbitrarily chosen positive real number. Two main cases are then considered:

Case 1: $z_0 \leqslant z < c\sqrt{\ln(\sqrt{n}/\,\mathsf{E}\,|X_1|^3)}$ ("moderate deviations");
Case 2: $z \geqslant z_0 \vee c\sqrt{\ln(\sqrt{n}/\,\mathsf{E}\,|X_1|^3)}$ ("large deviations");

here c is a positive constant.

In Case 1, of moderate deviations, the exponential tilting is performed, which may be presented as follows. Take some real $h > 0$ and let $\tilde{X}_1 = \tilde{X}_1^{(h,y)}, \dots, \tilde{X}_n = \tilde{X}_n^{(h,y)}$ be any r.v.'s such that

$$\mathsf{E}\, g(\tilde{X}_1, \dots, \tilde{X}_n) = \frac{\mathsf{E}\, e^{hS^{(y)}} g(X_1^{(y)}, \dots, X_n^{(y)})}{\mathsf{E}\, e^{hS^{(y)}}} \tag{3.2}$$

for all bounded (or for all nonnegative) Borel-measurable functions $g\colon \mathbb{R}^n \to \mathbb{R}$. Equivalently, one may require condition (3.2) only for Borel-measurable indicator functions g; clearly, such r.v.'s $\tilde{X}_i$ do exist. It is also clear that the r.v.'s $\tilde{X}_i$ are independent. These r.v.'s, the $\tilde{X}_i$'s, may be referred to as the tilted or, more specifically, h-tilted versions of the $X_i^{(y)}$'s. Clearly, without the truncation, the tilted versions of the original r.v.'s X_i may not exist, since $\mathsf{E}\, e^{hS}$ may be infinite even if $\mathsf{E}\,|X_i|^3 < \infty$ for all i. Using (3.2) with $g(x_1, \dots, x_n) = e^{-h(x_1+\cdots+x_n)}\,\mathrm{I}\{x_1 + \cdots + x_n > x\}$, it is easy to see that

$$\mathsf{P}(S^{(y)} > x) = \mathsf{E}\, e^{hS^{(y)}} \int_x^\infty \mathrm{d}u\, he^{-hu}\, \mathsf{P}(x < \tilde{S} \leqslant u) \tag{3.3}$$

for all real x, where $\tilde{S} := \tilde{X}_1 + \cdots + \tilde{X}_n$. Similarly, one can write

$$\mathsf{P}(BZ > x) = \mathsf{E}\, e^{hBZ} \int_x^\infty \mathrm{d}u\, he^{-hu}\, \mathsf{P}(x < BZ + B^2h \leqslant u) \tag{3.4}$$

for all real x, since any h-tilted version of the r.v. BZ has the distribution $N(B^2h, B^2)$. At that, good choices for y and h are of the form αx and $\eta x/B^2$, for some real parameters α and η in $(0,1)$.

So, to bound $|\,\mathsf{P}(S^{(y)} > z\sqrt{n}) - \mathsf{P}(Z > z)|$ (cf. (3.1)), one can demonstrate sufficient closeness of the terms $\mathsf{E}\, e^{hS}$ and $\mathsf{P}(x < \tilde{S} \leqslant u)$ in (3.3) to the corresponding terms $\mathsf{E}\, e^{hBZ}$ and $\mathsf{P}(x < BZ + B^2h \leqslant u)$ in (3.4). For each i, one notices that

$$\mathsf{E}\,|X_i^{(y)}|^3 e^{hX_i^{(y)}} \leqslant e^{hy}\,\mathsf{E}\,|X_i|^3 \tag{3.5}$$

and then shows that $\mathsf{E}\, e^{hX_i^{(y)}}$ is close enough to 1 and, somewhat more precisely, to $\mathsf{E}\, e^{hZ\sqrt{\mathsf{E}\, X_i^2}}$, and that the mean and variance of $\tilde{X}_i$ are close enough to $h\,\mathsf{E}\, X_i^2$ and $\mathsf{E}\, X_i^2$, respectively. So, one shows that $\mathsf{E}\, e^{hS^{(y)}}$ is close to $\mathsf{E}\, e^{hBZ}$, and the first two moments of $\tilde{S}^{(y)}$ are close enough to those of $BZ + B^2h$. Using now a uniform BE bound as in (1.1) – but for the $\tilde{X}_i$'s rather than the X_i's, one shows that $\mathsf{P}(x < \tilde{S} \leqslant u)$ is close enough to $\mathsf{P}(x < BZ + B^2h \leqslant u)$.

In Case 2, of large deviations, instead of the exponential tilting and a uniform BE bound, one employs an exponential inequality to bound $\mathsf{P}(S^{(y)} > x)$ and hence $\mathsf{P}(S^{(y)} > x) - \mathsf{P}(BZ > x)$ from above; for the lower bound on the latter difference, one simply uses $-\,\mathsf{P}(BZ > x)$.

3.1. A historical sketch of the problem of nonuniform BE bounds

The constant factors c_{nu} in the mentioned papers [19, 2] were not explicit. All papers known to this author with explicit values of c_{nu} followed the scheme of proof given by Nagaev [19], as delineated above.

Apparently the first such explicit value of c_{nu} was greater than 1955, as reported by Paditz [24]. In his dissertation [23], a much better value, 114.7, was presented, according to [21]; this author has not been able to access [23].

Michel [15] showed that in the iid case $c_{\mathsf{nu}} \leqslant c_u + 8(1 + e)$, which would be less than 30.2211, assuming the mentioned value 0.4748 for c_{u}, obtained in the later paper by Shevtsova [39].

Tysiak [43] stated c_{nu} to be no greater than 32.88. However, as was pointed out in [22], the expression $g_3(x)$ defined in [43, page 42] is missing a factor equal x^8, and this expression is used with $x > 3.4172$, so that the missing factor is greater than $3.4172^8 > 18593$. Even though this factor is missing only in one of the 5 summands comprising the final bound in [43], the overall impact of this omission could still be very large.

Mirakhmedov [16] claimed that Michel's constant, $c_{\mathsf{u}} + 8(1 + e)$, remains valid even without the iid assumption. However, as was pointed out by Wolf in his review of [16] (MR0780094), there was a mistake in [16], which used the Lyapunov ratio for the original r.v.'s X_i instead of that for the tilted truncated ones, $\tilde{X}_i$. There appear to be a number of other mistakes and gaps in [16] as well. In particular, it appears that the term $h^2B_n^2/2$ should not be on the right-hand side of [16, (10)]. Also, when using [16, (10)] to get [16, (12)], it is unclear how the term $\theta\,0.1876$ from [16, (10)] disappears in [16, (12)], given that L_n can be arbitrarily small. It is also unclear how one could get [16, (11)] from [16, (7) and (8)]. The subsequent letter to the editors [22] claimed that one could use the methods of the mentioned papers by Tysiak [43] and Mirakhmedov [16] to show that $c_{\mathsf{nu}} \leqslant 32.1530$. However, all the details for this claim in [22] are given in just four lines, and the displayed inequality there is incorrect, apparently due to a typo and mistaken editing (as the comparison with the manuscript of the letter shows).

Later, Paditz [25] claimed that $c_{\mathsf{nu}} < 31.935$. However, the proof in [25] contains a gap too. Namely, the case $c_{n,\delta,\alpha,\beta}(x) < K^2 < x^2$ is not considered in [25], where $c_{n,\delta,\alpha,\beta}(x)$ is a certain expression defined there. It appears likely that, when this missing case is considered, the value of the constant factor will go up. In the iid case, Gavrilenko [9] claimed $\big($instead of the bound of the form $c_{\mathsf{nu}}L/(1 + z^3)\big)$ the nonuniform bound $22.7707(L + 1/\sqrt{n})/(1 + z^3) \leqslant 2 \times 22.7707L/(1 + z^3)$. However, the proof in [9] has the same gap as the proof in [25]; also, the proof in [9] relies on the mentioned work [43]. Again in the iid case, Nefedova and Shevtsova [20] briefly stated that they had gone along the lines of the proof in [25] except using a better value for c_{u} (namely, 0.4784, obtained in [13]) in place of such a value (namely, 0.7915 [41]) used in [25], to get 25.80 for c_{nu}; it is unclear if they fixed the gap in [25] or even noticed it.

Finally, once again in the iid case, Nefedova and Shevtsova claimed in [21] that $c_{\mathsf{nu}} < 18.2$. However, there appears to be an error there as well. Namely, the first inequality in [21, (14)] is equivalent to the reverse of the last inequality on page 75 there, which latter is in turn equivalent to the condition $x^2 \geqslant c_n(x; \delta, a, b, c)$ in [21, Theorem 1], which is also equivalent to the second display on [21, page 75]; the expression $c_n(x; \delta, a, b, c)$ is defined in the first display on page 70 of [21]. So, for any given X, n, b, c, δ satisfying all the conditions of [21, Theorem 1], the first inequality in [21, (14)] and the last inequality on [21, page 75] can both hold only for one value of a. This wrong inequality in [21, (14)] is also used for [21, (16)].

This leaves, for now, 114.7 as the best (possibly correctly) established nonuniform BE constant factor c_{nu} – in the general, non-iid case. On the other hand, it follows from a result by Chistyakov [5, Corollary 1] that c_{nu} is necessarily no less than 1, and this lower bound on c_{nu} is asymptotically exact in a certain sense for $z \to \infty$. Apparently, this has been the best known lower bound on c_{nu}. However, it is easy to improve this bound slightly and show that necessarily

$$c_{\mathsf{nu}} > 1.0135; \tag{3.6}$$

this can be done by letting X_1 have the centered Bernoulli distribution with parameter $p = 8/100$ and then letting $n = 1$ and $z \uparrow 1 - p$. However, it was shown by Bentkus [1], the best constant factor for $n = 1$ will be 1 if $1 + z^3$ in (3.1) is replaced by z^3; it is also conjectured in [1] that the same constant factor, 1, will be good for all n.

Thus, in the non-iid case the apparently best known lower bound on c_{nu} is over 100 times smaller than the best established upper bound on c_{nu}, and this gap is about 30 times in the iid case. So, the situation with c_{nu} is much worse than that with c_{u}.

3.2. Possible improvements of Nagaev's method

A crucial component of the mentioned method offered by Nagaev [19] and used in the subsequent papers [2, 24, 23, 15, 43, 16, 22, 25, 9, 20, 21] is an exponential inequality. However, the exponential bounds used in all of those papers are not the best possible ones. An optimal exponential bound, in terms of the first two moments and truncated absolute third moments of the X_i's was given by Pinelis and Utev [37]. In fact, the paper [37] provided a general method to obtain optimal exponential bounds, along with a number of specific applications of the general method.

However, even the best possible *exponential* bounds, say for sums of independent r.v.'s, can be significantly improved. The reason for this is that the class of exponential moments functions is very small (even though analytically very simple to deal with). Using a much richer class of moments functions, Pinelis [34] obtained the following result. Let $X_1, \dots, X_n$ be independent random variables (r.v.'s), with the sum $S := X_1 + \cdots + X_n$. For any $a > 0$ and $\theta > 0$, let Γ_{a^2} and Π_θ stand for any independent r.v.'s such that Γ_{a^2} has the normal distribution with parameters 0 and a^2, and Π_θ has the Poisson distribution with parameter θ. Let also $\tilde\Pi_\theta := \Pi_\theta - \mathsf{E}\,\Pi_\theta = \Pi_\theta - \theta$. Let σ, y, and β be any positive real numbers such that $\varepsilon := \frac{\beta}{\sigma^2 y} \in (0,1)$. Suppose that $\sum_i \mathsf{E}\, X_i^2 \leqslant \sigma^2$, $\sum_i \mathsf{E}(X_i)_+^3 \leqslant \beta$, $\mathsf{E}\, X_i \leqslant 0$, and $X_i \leqslant y$, for all i. Let $\eta_{\varepsilon,\sigma,y} := \Gamma_{(1-\varepsilon)\sigma^2} + y\tilde\Pi_{\varepsilon\sigma^2/y^2}$. Then it is proved in [34] that

$$\mathsf{E}\, f(S) \leqslant \mathsf{E}\, f(\eta_{\varepsilon,\sigma,y}) \tag{3.7}$$

for all twice continuously differentiable functions f such that f and f'' are nondecreasing and convex. A corollary of this result is that for all $x \in \mathbb{R}$

$$\mathsf{P}(S \geqslant x) \leqslant \inf_{t\in(-\infty,x)} \frac{\mathsf{E}(\eta_{\varepsilon,\sigma,y} - t)_+^3}{(x-t)^3} \leqslant c_{3,0}\, \mathsf{P}^{\mathsf{LC}}(\eta_{\varepsilon,\sigma,y} \geqslant x), \tag{3.8}$$

where $c_{3,0} := \frac{2e^3}{9} \approx 4.46$ and the function $\mathbb{R} \ni x \mapsto \mathsf{P}^{\mathsf{LC}}(\eta \geqslant x)$ is defined as the least log-concave majorant over $\mathbb{R}$ of the tail function $\mathbb{R} \ni x \mapsto \mathsf{P}(\eta \geqslant x)$ of a r.v. η. The bounds in (3.7) and (3.8) are much better than even the best exponential bounds (expressed in the same terms).

A trade-off here is that the bounds given in (3.8) are significantly more difficult to deal with, especially analytically, than exponential bounds. However, this can be done, as shown in the following discussion. In accordance with what was pointed out above, one needs an exponential bound (or a better one) only in Case 2, of large deviations, when $z \geqslant z_0 \vee c\sqrt{\ln(\sqrt{n}/\,\mathsf{E}\,|X_1|^3)}$, which implies

$$\mathsf{E}\,|X_1|^3/\sqrt{n} \geqslant e^{-z^2/c^2}. \tag{3.9}$$

Also, by (3.8), for any real $\tau < 1$

$$(1-\tau)^3\,\mathsf{P}(S^{(y)} > x) \leqslant \sum_{j=0}^{\infty} Q_j \frac{\lambda^j}{j!} e^{-\lambda},$$

where

$$x := Bz = z\sqrt{n}, \quad y = \alpha x, \quad \alpha \in (0,1), \quad Q_j := \Big(\frac{\alpha_1}{z}\Big)^3\,\mathsf{E}(Z+u_j)_+^3,$$
$$\alpha_1 := \sqrt{1-az_0^2/\alpha}, \quad a := \frac{\mathsf{E}(y\wedge (X_1)_+)^3}{z^3\sqrt{n}}, \quad u_j := \alpha\Big(j - \frac{\tau}{\alpha} - \lambda\Big)z, \quad \lambda := \frac{a}{\alpha^3}. \tag{3.10}$$

Assume now that $\tau c^2 \geqslant 2$, where c is as in (3.9). Since $z \geqslant z_0$, one has

$$Q_0 \leqslant C_0 \frac{\mathsf{E}\,|X_1|^3}{z^3\sqrt{n}}, \tag{3.11}$$

where

$$C_0 := e^{z_0^2/c^2}\,\mathsf{E}(Z-\tau z_0)_+^3; \tag{3.12}$$

here one uses the fact that $e^{\beta t^2}\,\mathsf{E}(Z-t)_+^3$ is decreasing in $t \geqslant 0$ provided that $\beta \leqslant 1/2$; in fact, this decrease is fast, especially when $\beta < 1/2$. Note also that $\mathsf{E}(Z-t)_+^3 = \big(t^2+2\big)\,\varphi(t) - t\,\big(t^2+3\big)\,\bar{\Phi}(t)$ for all real t, where φ and $\bar{\Phi}$ are the density and tail functions of Z.

Next, since $\mathsf{E}\,g(\beta Z)$ is nondecreasing in $\beta \geqslant 0$ for any convex function g,

$$Q_j \leqslant (\alpha_1/z_0)^3\,\mathsf{E}(Z+u_{j0})_+^3, \quad \text{where} \quad u_{j0} := \alpha(j-\tau/\alpha-\lambda)z_0.$$

Using now the identity $\mathsf{E}(Z+t)_+^3 = t^3+3t+\mathsf{E}(Z-t)_+^3$ for all real t and the decrease of $\mathsf{E}(Z-t)_+^3$ in $t \in \mathbb{R}$, one has

$$\sum_1^{\infty} Q_j \frac{\lambda^j}{j!} e^{-\lambda} \leqslant C_1(a) \frac{\mathsf{E}\,|X_1|^3}{z^3\sqrt{n}}, \tag{3.13}$$

where

$$C_1(a) := \Big(\frac{\alpha_1}{\alpha z_0}\Big)^3 \Big[\sum_1^{\infty}(u_{j0}^3+3u_{j0})\frac{\lambda^{j-1}}{j!} + \mathsf{E}(Z-u_{10})_+^3 + \mathsf{E}(Z-u_{20})_+^3\frac{\lambda^{2-1}}{2!}e^{\lambda}\Big]e^{-\lambda},$$

with λ as defined in (3.10). The sum $\sum_1^\infty$ in the above expression of $C_1(a)$ is easy to evaluate explicitly. Also, since the left-hand side of (3.1) can never exceed 1, without loss of generality

$$a \leqslant a_{\max}, \tag{3.14}$$

with $a_{\max} = 1/c_{\mathsf{nu}}$; working a bit harder, one may assume (3.14) with $a_{\max}$ significantly smaller than $1/c_{\mathsf{nu}}$. Next, it appears that for values of the parameters α and τ that have a chance to be optimal or quasi-optimal, the factor $C_1(a)$ will be decreasing in $a \in [0, a_{\max}]$. Therefore and in view of (3.11) and (3.13), one will have

$$\mathsf{P}(S^{(y)} > x) \leqslant \frac{C_0 + C_1(0+)}{(1-\tau)^3}\,\frac{\mathsf{E}\,|X_1|^3}{z^3\sqrt{n}}, \tag{3.15}$$

with C_0 as in (3.12) and $C_1(0+) = (\alpha z_0)^{-3}\,\mathsf{E}(Z+(\alpha-\tau)z_0)_+^3$. One can improve the above estimates by partitioning the interval $[0, a_{\max}]$ into a number of smaller subintervals and then considering the corresponding cases depending on which of the subintervals the value of a is in.

Thus, it is shown that $\mathsf{P}(S^{(y)} > x)$ can be appropriately bounded using the better-than-exponential bound in (3.8), and at that in a rather natural manner and incurring almost no losses. It should be clear that the expression on the right-hand side of inequality (3.15) will become a term in a bigger expression that is an upper bound on the left-hand side of (3.1). That latter, bigger expression will then have to be (quasi-)minimized with respect to z_0, α, η, τ, and the other parameters, subject to the necessary restrictions on their values.

Also, one can use ideas from [17, 18, 35, 36] to improve the estimation of the effect of truncation, as compared with the way that was done in the mentioned papers [19, 2, 24, 23, 15, 43, 16, 22, 25, 9, 20, 21], as well as more "synthetic" ways to bound moments of the tilted distribution – cf. results in [30, 29, 33, 32]. In addition, as in [21], one can use the uniform bound $0.3328(\mathsf{E}\,|X_1|^3 + 0.429)/\sqrt{n}$ from [39], which is smaller than the previously mentioned bound of the classical form $c_{\mathsf{u}}L = c_u\,\mathsf{E}\,|X_1|^3/\sqrt{n}$ with $c_{\mathsf{u}} = 0.4748$. There are a few other potentially useful modifications. Thus, the improvements concern every one of the three major ingredients of Nagaev's method listed beginning on page 3. By utilizing the above ideas, one may hope to improve the upper bound on c_{nu} to about 10 in the iid case and to about 12 in the general case. When and if such an objective is attained, the gap between the available upper and lower bounds on c_{nu} will be decreased, at least, about 3 times in the iid case and about 10 times in the general case.

However, significant further progress after that seems unlikely within the framework of the method of [19]. One of the main obstacles here is the factor e^{hy} as in (3.5). Since good choices for y and h turn out to be αx and $\eta x/B^2$ with α and η somewhat close to 0.5, this factor will then be something like $e^{z^2/4}$, which is large for large enough z.

Yet, the factor e^{hy} is the best possible one in (3.5) (even assuming that $X_i^{(y)} = X_i$ and hence $X_i^{(y)}$ is zero-mean). Such a large factor is necessary when X_i has a two-point distribution highly skewed to the right. On the other hand, certain considerations suggest that the least favorable situation in Case 1 of moderate deviations is when n is very large but z is not so, and then the mentioned least favorable distribution (for the uniform BE bound) given by (1.3) is only slightly skewed. This creates a significant tension in using the exponential tilt.

One may try to reduce the factor e^{hy} by decreasing α and hence y – but this will increase fast the effect of truncation, which is (at least roughly) proportional to $1/\alpha^3$.

Even if one were able to get rid of the factor e^{hy} altogether, the corresponding uniform BE bound on the rate of convergence to the probability $\mathsf{P}(x < BZ + B^2h \leqslant u)$ in (3.4) would still seem relatively too large, since this probability itself is less than $\mathsf{P}(x < BZ + B^2h) = \mathsf{P}\big(Z > (1-\eta)z\big)$ and therefore is rather small for what appears to be the least favorable values of z, such as 2.5 to 3.5 (and values of η typically not too far from 0.5). In contrast, the mentioned asymptotic lower bound by Esseen [8] (recall (1.3)) is attained for z close to 0; furthermore, the corresponding asymptotic expression is rather highly peaked near the maximum and is thus much smaller outside of a neighborhood of the maximum point.

Yet another apparently powerful cause of tension is as follows. After the X_i's have been truncated, a natural bound on $|\,\mathsf{P}(S^{(y)} > x) - \mathsf{P}(BZ > x)|$, obtained via either the exponential tilt or a Stein-type method, decays in an exponential rather than power fashion; see e.g. the results [4] and [28], which imply an upper bound of the form $c(\lambda)\frac{\mathsf{E}\,|X_1|^3}{e^{\lambda z}\sqrt{n}}$ for real $\lambda > 0$, say in the iid case. The factor $1/e^{\lambda z}$ decays much faster than $1/(1+z^3)$ when z is large. However, the former factor may be much greater than the latter, especially if λ is not large and z is not very large. For instance, if $\lambda = 1/2$ as in [4], then $\max_{z>0} \frac{1}{e^{\lambda z}}\big/\frac{1}{1+z^3} = 10.8\dots$, attained at $z = 5.9719\dots$.

In the next section, a new approach to obtaining nonuniform BE bounds is described, based on the Fourier method, complemented by extremal problem methods.

4. A new way to obtain nonuniform BE bounds

Take any function $h \in C^1$ such that (the limit) $\mathfrak{G}(h)$ exists (and is) in $\mathbb{R}$ and $h(t)/t \to 0$ as $|t| \to \infty$; here and in what follows, C^k denotes the class of all k times continuously differentiable complex-valued functions defined on $\mathbb{R}$. Take any real $x \in \mathbb{R}$. Note that $\mathfrak{G}(1)(x) = \frac{1}{2}\operatorname{sign} x$ – say, by (2.6) with $X = 0$. So, writing $\mathfrak{G}(h) = h(0)\mathfrak{G}(1) + \mathfrak{G}\big(h - h(0)\big)$ and evaluating the p.v.-integral in the expression for $\mathfrak{G}\big(h - h(0)\big)$ by parts, one has

$$x\mathfrak{G}(h)(x) = \tfrac{1}{2}\, h(0)\, x \operatorname{sign} x + i\, \mathfrak{G}(\Lambda h)(x) \tag{4.1}$$

if $x \neq 0$, where the linear operator Λ is defined by the formula

$$(\Lambda h)(t) := -t\,\frac{\mathrm{d}}{\mathrm{d}t}\,\frac{h(t) - h(0)}{t} = -\frac{h(0) - [h(t) + h'(t)(-t)]}{t} \tag{4.2}$$

for $t \neq 0$. In fact, identity (4.1) holds for $x = 0$ as well, in view of the definitions of $\mathfrak{G}$ and Λ. By induction, for all $k \in \mathbb{N} := \{1, 2, \dots\}$

$$(\Lambda^k h)(t) = -k!\, t^{-k}\Big(h(0) - \sum_{j=0}^{k} h^{(j)}(t)\,\frac{(-t)^j}{j!}\Big) = (-1)^k \int_0^1 [h^{(k)}(t) - h^{(k)}(\alpha t)]\, k\alpha^{k-1}\,\mathrm{d}\alpha \tag{4.3}$$

if $h \in C^k$ and $t \neq 0$, and hence $(\Lambda^k h)(0) = 0$. So, iterating (4.1), one has

$$\mathfrak{G}(h)(x) = \frac{h(0)\operatorname{sign} x}{2} + \Big(\frac{i}{x}\Big)^k \mathfrak{G}(\Lambda^k h)(x) \tag{4.4}$$

for all real $x \neq 0$, all $k \in \mathbb{N}$, and functions h such that

$$h \in C^k, \quad \mathfrak{G}(h) \text{ exists in } \mathbb{R}, \quad \text{and} \quad h^{(j)}(t)/t \to 0 \text{ for all } j = 1, \dots k \text{ as } |t| \to \infty. \tag{4.5}$$

More generally,

$$\mathfrak{G}(h)(x) = \frac{h(0)\operatorname{sign} x}{2} + \frac{i}{2\pi}\sum_{j=1}^{k}\frac{h^{(j)}(0+) - h^{(j)}(0-)}{j\,(ix)^j} + \Big(\frac{i}{x}\Big)^k \mathfrak{G}(\Lambda^k h)(x) \tag{4.6}$$

for all real $x \neq 0$, all $k \in \mathbb{N}$, and all functions h such that

$$\begin{gathered} h \in C(\mathbb{R}), \quad h \in C^k(\mathbb{R}\backslash\{0\}), \quad \mathfrak{G}(h) \text{ exists in } \mathbb{R}, \quad \text{and} \\ \text{for each } j \in \{1, \dots k\} \text{ there exists } h^{(j)}(0\pm) \in \mathbb{R} \text{ and } h^{(j)}(t)/t \to 0 \text{ as } |t| \to \infty. \end{gathered} \tag{4.7}$$

The condition $h \in C^k(\mathbb{R}\backslash\{0\})$ in (4.7) can be slightly relaxed, to the following:

$$h \in C^{k-1}(\mathbb{R}\backslash\{0\}) \quad \text{and} \quad h^{(k-1)} \text{ is of locally bounded variation on } \mathbb{R}\backslash\{0\}, \tag{4.8}$$

with $\mathfrak{G}(\Lambda^k h)(x)$ then understood as $\mathfrak{G}(\widetilde{\Lambda^k} h)(x) + (-1)^k \tilde{\mathfrak{G}}(h^{(k-1)})(x)$, where $(\widetilde{\Lambda^k} h)(t) := -k!\, t^{-k}\big(h(0) - \sum_{j=0}^{k-1} h^{(j)}(t)\,\frac{(-t)^j}{j!}\big)$ for $t \neq 0$ and $\tilde{\mathfrak{G}}(h^{(k-1)})(x) := \frac{i}{2\pi}\,\text{p.v.}\int_{-\infty}^{\infty} e^{-itx}\,\frac{\mathrm{d}h^{(k-1)}(t)}{t}$.

Identity (4.4) immediately implies

Theorem 4.1. *Take any $k \in \mathbb{N}$ and any real $T > 0$ and $x \geqslant 0$. Let X be any r.v. with $\mathsf{E}\,|X|^k < \infty$. Let f denote the c.f. of X. Let M be as in (2.2), with the additional requirement that $M \in C^k$. Then*

$$-\,i^k \mathfrak{G}\big(\Lambda^k\, r_{T,-}\big)(x) \leqslant x^k\,\mathsf{P}(X > x) \leqslant x^k\,\mathsf{P}(X \geqslant x) \leqslant -i^k \mathfrak{G}\big(\Lambda^k\, r_{T,+}\big)(x), \tag{4.9}$$

where

$$r_{T,\pm}(\#) := M_T(\mp\#) f(\#). \tag{4.10}$$

Remark. Condition $\mathsf{E}\,|X|^k < \infty$ in Theorem 4.1 implies $f \in C^k$, so that (4.5) holds with $g = r_{T,\pm}$.

As was mentioned, the Prawitz smoothing filter M given by (2.7) provides the tightest, in a certain sense, upper and lower bounds in (2.2) on the d.f. of X. However, it is not smooth enough to be used in Theorem 4.1 in the most interesting in applications case $k = 3$. Namely, that M is not even in C^1 – whereas one needs $M \in C^3$ in Theorem 4.1 for $k = 3$.

There are a number of ways to develop such a smooth enough smoothing filter. Some of them can be based on Proposition 5.1 in Section 5 of this paper; see e.g. the function $M = M_{0,2}$ given by formula (5.4).

The identity (2.6) can be rewritten in the following more general and hence sometimes more convenient form.

Proposition 4.2. *Let L be any complex-valued function of bounded variation on $\mathbb{R}$, and let ℓ be its Fourier–Stieltjes transform, so that $\ell(t) = \int_{-\infty}^{\infty} e^{itx}\, \mathrm{d}L(x)$ for all real t. Assume also that L is regularized so that $2L(x) = L(x-) + L(x+)$ for all $x \in \mathbb{R}$ and extended to $[-\infty, \infty]$ so that $L(\pm\infty) = \lim_{x\to\pm\infty} L(x)$. Then*

$$L(x) - \tfrac{1}{2}\,[L(\infty) - L(-\infty)] = \mathfrak{G}(\ell)(x) \quad \textit{for all real } x. \tag{4.11}$$

This follows immediately from (2.6), because (i) both sides of (4.11) are linear in L and (ii) any regularized function of bounded variation on $\mathbb{R}$ is a linear combination (with complex coefficients) of regularized distribution functions.

Suppose that $\ell\colon \mathbb{R} \to \mathbb{C}$ is a function which may depend on a number of parameters. For brevity, let us say that the function ℓ is a *quasi-c.f.* if it can be represented as a linear combination of k c.f.'s with (possibly complex) coefficients such that the length k of the combination and the coefficients are bounded uniformly over all possible values of the parameters.

Clearly, the product of two quasi-c.f.'s is a quasi-c.f. Also, any linear combination of two quasi-c.f.'s is a quasi-c.f., provided that the coefficients of the combination are bounded uniformly over all possible values of the parameters. Moreover, one has the following simple proposition.

Proposition 4.3. *Take any natural m. Let N denote the c.f. of a r.v. Y whose distribution may depend on a number of parameters. Suppose that $\mathsf{E}\,|Y|^m$ is (finite and) bounded uniformly over all possible values of the parameters. Then the mth derivative $N^{(m)}$ of N is a quasi-c.f.*

Proof of Proposition 4.3. Let us exclude the trivial case when $\mathsf{E}\,|Y|^m = 0$. If m is even or $\mathsf{E}\,Y_+^m = 0$ or $\mathsf{E}\,Y_-^m = 0$, then $\widetilde{N^{(m)}}(\#) := \frac{\mathsf{E}\,Y^m e^{iY\#}}{\mathsf{E}\,Y^m}$ is a c.f., and $N^{(m)} = (i^m\,\mathsf{E}\,Y^m)\widetilde{N^{(m)}}$, so that $N^{(m)}$ is a quasi-c.f. In the remaining case one has $\mathsf{E}\,Y_+^m > 0$ and $\mathsf{E}\,Y_-^m > 0$, so that one can similarly write $N^{(m)}(\#) = i^m\,\mathsf{E}\,Y_+^m\,\frac{\mathsf{E}\,Y_+^m e^{iY\#}}{\mathsf{E}\,Y_+^m} + (-i)^m\,\mathsf{E}\,Y_-^m \frac{\mathsf{E}\,Y_-^m e^{iY\#}}{\mathsf{E}\,Y_-^m}$. □

A quick proof of Nagaev's nonuniform BE bound (3.1) can be easily obtained based on Theorem 4.1. Indeed, let $T = c_T\sqrt{n}/\beta_3$, where $\beta_3 := \mathsf{E}\,|X_1|^3$ and c_T is a small enough positive real constant. Let $A \underset{\frown}{<} B$ mean $|A| \leqslant CB$ for some absolute constant C. Let $X := S/\sqrt{n}$. If $T \leqslant 1$ then $1 \underset{\frown}{<} \frac{\beta_3}{\sqrt{n}}$. So, for all real $x \geqslant 0$, by the Markov and Rosenthal inequalities, $(1+x^3)\,\mathsf{P}(X \geqslant x) \leqslant 1 + \mathsf{E}\,|X|^3 \underset{\frown}{<} 1 + \frac{\beta_3}{\sqrt{n}} \underset{\frown}{<} \frac{\beta_3}{\sqrt{n}}$ and similarly $(1+x^3)\,\mathsf{P}(Z \geqslant x) \underset{\frown}{<} \frac{\beta_3}{\sqrt{n}}$, whence (3.1) follows.

It remains to consider the case $T > 1$. Note that then $n > (\beta_3/c_T)^2 \geqslant 3$ and hence $n \geqslant 4$ provided that $c_T \leqslant 1/\sqrt{3}$.

In view of the uniform BE bound, Theorem 4.1, and (4.3), in order to prove (3.1) it is enough to show that $\mathfrak{G}\big(r'''_{1,f}(\alpha\#) - r'''_{1,g}(\alpha\#)\big) \underset{\frown}{<} \beta_3/\sqrt{n}$ and $\mathfrak{G}\big(r'''_{2,f}(\alpha\#)\big) \underset{\frown}{<} \beta_3/\sqrt{n}$ over $\alpha \in (0,1]$, where

$r_{j,f}(\#) := M_j(\frac{\#}{T})f(\#)$, $j = 1, 2$, the M_j's are as in (2.5), M is (say) as in (5.4), f is the c.f. of $X := S/\sqrt{n}$, and $g(\#) := e^{-\#^2/2}$ (so that g may be considered as a special case of f). One has

$$r'''_{j,f}(\#) = \sum_{q=0}^{3} \binom{3}{q} \frac{1}{T^q} M_j^{(q)}\Big(\frac{\#}{T}\Big) f^{(3-q)}(\#). \tag{4.12}$$

By (5.4) and (5.6), M_1 is the c.f. of a distribution with a finite 4th moment, whereas $M_2 = \kappa M_1'$. Hence, by Proposition 4.3, $M_j^{(q)}$ is a quasi-c.f. for each pair $(j, q) \in \{1, 2\} \times \{0, 1, 2, 3\}$, and then so is $M_j^{(q)}(\frac{\#}{T})$. Similarly, f is the c.f. of the r.v. X with $\mathsf{E}\,|X|^3 \lesssim 1 + \frac{\beta_3}{\sqrt{n}} \lesssim 1$, by the Rosenthal inequality and the case condition $T > 1$. So, again by Proposition 4.3, $f^{(3-q)}$ is a quasi-c.f. for each $q \in \{0, 1, 2, 3\}$. Thus, $M_j^{(q)}(\frac{\alpha\#}{T}) f^{(3-q)}(\alpha\#)$ is a quasi-c.f. and, by Proposition 4.2, $\mathfrak{G}\big(M_j^{(q)}(\frac{\alpha\#}{T}) f^{(3-q)}(\alpha\#)\big) \lesssim 1$, for each $(j, q) \in \{1, 2\} \times \{0, 1, 2, 3\}$. Therefore and because $T > 1$,

$$\mathfrak{G}\Big(\binom{3}{q} \frac{1}{T^q} M_j^{(q)}(\tfrac{\alpha\#}{T}) f^{(3-q)}(\alpha\#)\Big) \lesssim \frac{1}{T^q} \leqslant \frac{1}{T} \lesssim \frac{\beta_3}{\sqrt{n}} \quad \text{for each } (j, q) \in \{1, 2\} \times \{1, 2, 3\}; \tag{4.13}$$

note that $q = 0$ is not included here.

It remain to show that $\mathfrak{G}_{1\alpha}(f''' - g''') \lesssim \frac{\beta_3}{\sqrt{n}}$ and $\mathfrak{G}_{2\alpha}(f''') \lesssim \frac{\beta_3}{\sqrt{n}}$, where

$$\mathfrak{G}_{j\alpha}(h)(x) := \mathfrak{G}\big(M_j(\tfrac{\alpha\#}{T})(h)(\alpha\#)\big)(x). \tag{4.14}$$

For $j \in \{0, 1, 2, 3\}$, introduce $f_1^{(j)}(t) := \big(\frac{\mathrm{d}}{\mathrm{d}t}\big)^j f_1(t)$ and $f_{1n}^{(j)}(t) := f_1^{(j)}(t/\sqrt{n})$, where f_1 denotes the c.f. of X_1. Similarly, starting with $g_1 := g$ in place of f_1, define $g_{1n}^{(j)}$, and then let $d_{1n}^{(j)} := f_{1n}^{(j)} - g_{1n}^{(j)}$ and $h_{1n}^{[j]} := \big|f_{1n}^{(j)}\big| \vee \big|g_{1n}^{(j)}\big|$; omit superscripts ${}^{(0)}$ and ${}^{[0]}$. Note that $f = f_{1n}^n$ and hence $\sqrt{n} f''' = f_{31} + f_{32} + f_{33}$, where $f_{31} := (n-1)(n-2) f_{1n}^{n-3} \big(f_{1n}^{(1)}\big)^3$, $f_{32} := 3(n-1) f_{1n}^{n-2} f_{1n}^{(1)} f_{1n}^{(2)}$, and $f_{33} := f_{1n}^{n-1} f_{1n}^{(3)}$; do similarly with g and g_1 in place of f and f_1. By Proposition 4.3, $M_j(\frac{\alpha\#}{T}) f_{33}/\beta_3$ is a quasi-c.f. and hence, by Proposition 4.2, $\mathfrak{G}_{j\alpha}(f_{33}) \lesssim \beta_3$, for $j \in \{1, 2\}$.

So, it suffices to show that $\mathfrak{G}_{1\alpha}(f_{3k} - g_{3k}) \lesssim \beta_3$ and $\mathfrak{G}_{2\alpha}(f_{3k}) \lesssim \beta_3$ for $k \in \{1, 2\}$. This can be done in a straightforward manner using the following estimates for $j \in \{0, 1, 2, 3\}$ and $|t| \leqslant T$: $M_1 \lesssim 1$, $M_2(\frac{t}{T}) \lesssim \frac{|t|}{T} \lesssim |t|\beta_3/\sqrt{n}$, $h_{1n}(t)^{n-j} \leqslant e^{-ct^2}$ (where c is a positive real number depending only on the choice of c_T), $h_{1n}^{[1]}(t) \lesssim |t|/\sqrt{n}$, $h_{1n}^{[2]}(t) \lesssim 1$, $|d_{1n}^{(j)}(t)| \lesssim \beta_3(|t|/\sqrt{n})^{3-j}$, and hence $f_{1n}^{n-j}(t) - g_{1n}^{n-j}(t) \lesssim |t|^3 e^{-ct^2} \beta_3/\sqrt{n}$; cf. e.g. [26, Ch. V, Lemma 1]. For instance, $|f_{31} - g_{31}| \lesssim n^2(D_{311} + D_{312})$, where $D_{311}(t) := \big(|f_{1n}^{n-3} - g_{1n}^{n-3}| \big(h_{1n}^{[1]}\big)^3\big)(t) \lesssim |t|^3 e^{-ct^2} \frac{\beta_3}{\sqrt{n}} \big(\frac{|t|}{\sqrt{n}}\big)^3$ and $D_{312}(t) := \big(h_{1n}^{n-3} \big(h_{1n}^{[1]}\big)^2 |d_{1n}^{(1)}|\big)(t) \lesssim e^{-ct^2} \big(\frac{|t|}{\sqrt{n}}\big)^2 \beta_3 \big(\frac{|t|}{\sqrt{n}}\big)^2$, so that $\mathfrak{G}_{1\alpha}(f_{31} - g_{31}) \lesssim \int_{-\infty}^{\infty} (t^6 + t^4) e^{-ct^2} \beta_3 \frac{\mathrm{d}t}{|t|} \lesssim \beta_3$. □

Of course, the above argument is rather crude and yet it demonstrates that the method based on the smoothing inequalities (4.9) is quite effective. It also strongly suggests that this method can be used further, in order to obtain an explicit and appropriately small upper bound on the constant factor c_{nu}.

Let us now discuss some of the refinements that could be used within the general framework of the above *quick proof of* (3.1).

There, in particular, we needed to bound

$$L(H) := \int_0^1 \big[\mathfrak{G}\big(H(\alpha\#)\big)(x) - \mathfrak{G}\big(H(\#)\big)\big]\, 3\alpha^2 \,\mathrm{d}\alpha, \tag{4.15}$$

where H is of the form $M_1(\frac{\#}{T})(f_{3k}-g_{3k})$ or $M_2(\frac{\#}{T})f_{3k}$ for $k \in \{1,2\}$ – recall (4.9), (4.10), and (4.3). Tacitly, that bounding was then done using the trivial inequalities

$$|L(H)| \leqslant 2 \sup_{\alpha\in(0,1]} \big|\mathfrak{G}\big(H(\alpha\#)\big)(x)\big| \leqslant 2\,\frac{1}{2\pi}\int_\infty^\infty |H(t)|\frac{\mathrm{d}t}{|t|}, \tag{4.16}$$

where in turn we used the definition (2.4) of $\mathfrak{G}$ and the trivial identity $|e^{-itx}| = 1$ for real t and x; the integral in (4.16) exists even in the Lebesgue sense, since $|f_{3k}(t) - g_{3k}(t)| = O(|t|)$ and $|M_2(t)| = O(|t|)$.

In fact, the factor 2 in the last bound in (4.16) on $|L(H)|$ can be removed, so that one have

$$|L(H)| \leqslant \frac{1}{2\pi}\int_\infty^\infty |H(t)|\frac{\mathrm{d}t}{|t|}. \tag{4.17}$$

Indeed, first of all note here that the factor α can be easily moved, in a way, from the argument of the general and hard to control function H into that of the much simpler and more specific exponential function, using the simple identity

$$\mathfrak{G}\big(H(\alpha\#)\big)(x) = \mathfrak{G}\big(H(\#)\big)(\tfrac{x}{\alpha}), \tag{4.18}$$

which implies that

$$L(H) = \frac{i}{2\pi}\int_\infty^\infty I(tx)H(t)\frac{\mathrm{d}t}{t}, \quad \text{where} \tag{4.19}$$

$$I(u) := \int_0^1 (e^{-iu/\alpha} - e^{-iu})\,3\alpha^2\,\mathrm{d}\alpha. \tag{4.20}$$

Now (4.17) follows immediately from

Proposition 4.4. *The expression $g(u) := |I(u)|^2$ is even in $u \in \mathbb{R}$ and (strictly) increases from 0 to 1 as $|u|$ increases from 0 to ∞; in particular, it follows that $|I(u)| \in [0,1)$ for all real u. Moreover, the function g has the following generalized concavity property: $-u^3\big(u^{-5}g'(u)\big)'$ is completely monotone in $u > 0$ (in Bernstein's sense – see e.g. [27, Chapter 2]); in particular, $g(v^{1/6})$ is concave in $v > 0$.*

Thus, the conclusion in Proposition 4.4 that $|I(u)| \in [0,1)$ for all real u can be seen as a rather sophisticated replacement for the trivial identity $|e^{-itx}| = 1$ for real t and x, which latter was used to obtain the rightmost bound in (4.16).

Moreover, one can easily obtain (and then use in (4.19)) an upper bound on $|I(u)|$ which is significantly less than 1 for small enough values of $|u|$. This can be done by closely bounding the values of $|I(u)|$ for a finite number of values of u and then using the monotonicity property of $|I|$ provided by Proposition 4.4.

Graphs of $\operatorname{Re} I$, $\operatorname{Im} I$, and $|I|$ over the interval $[-6\pi, 6\pi]$ are shown in Figure 1. It seems plausible that $g(u)$ is concave in $u > 0$; however, that probably would be hard to prove.

Proof of Proposition 4.4. Note that $I(-u) = \overline{I(u)}$ for all real u. So, the function $g = I\overline{I}$ is indeed even.

Take now any real $u > 0$. Integrating by parts and then changing the integration variable, one has

$$iu^{-3}I(u) = u^{-2}\int_0^1 e^{-iu/\alpha}\,\alpha\,\mathrm{d}\alpha = \mathcal{E}(u) := \mathcal{E}_3(u), \quad \text{where} \quad \mathcal{E}_j(u) := \int_u^\infty e^{-iz}\frac{\mathrm{d}z}{z^j} \tag{4.21}$$



Fig 1: Graphs of $\operatorname{Re} I$, $\operatorname{Im} I$, and $|I|$.

for $j > 0$. So, $g(u) = u^6\,\mathcal{E}(u)\overline{\mathcal{E}(u)}$,

$$\begin{aligned}
g_1(u) &:= \tfrac{1}{2}\,u^{-5}g'(u) = 3\mathcal{E}(u)\overline{\mathcal{E}(u)} - u^{-2}\operatorname{Re}\big(e^{iu}\mathcal{E}(u)\big),\\
u^3 g_1'(u) &= \frac{1}{u^2} - \operatorname{Re}\big((4+iu)e^{iu}\mathcal{E}(u)\big)\\
&= \frac{1}{u^2} - \operatorname{Re}\Big((4+iu)\int_u^\infty e^{-i(z-u)}\,\frac{\mathrm{d}z}{z^3}\Big)\\
&= \frac{1}{u^2} - \operatorname{Re}\Big((4+iu)\int_0^\infty e^{-iv}\,\frac{\mathrm{d}v}{(v+u)^3}\Big)\\
&= \int_0^\infty e^{-us}s\,\mathrm{d}s - \operatorname{Re}\Big((4+iu)\int_0^\infty e^{-iv}\,\mathrm{d}v\int_0^\infty \tfrac{1}{2}e^{-(v+u)s}s^2\,\mathrm{d}s\Big).
\end{aligned}$$

Noting now that $\int_0^\infty e^{-iv}e^{-vs}\,\mathrm{d}v = \frac{1}{s+i}$ and $\operatorname{Re}\frac{4+iu}{s+i} = \frac{4s+u}{s^2+1}$ for all real $s > 0$, and introducing $w_1(s) := 2\frac{s-s^3}{s^2+1}$ and $w_2(s) := \frac{-s^2}{s^2+1}$, write

$$\begin{aligned}
2u^3 g_1'(u) &= \int_0^\infty e^{-us}w_1(s)\,\mathrm{d}s + \int_0^\infty e^{-us}w_2(s)\,u\,\mathrm{d}s\\
&= \int_0^\infty e^{-us}w_1(s)\,\mathrm{d}s + \int_0^\infty e^{-us}w_2'(s)\,\mathrm{d}s = -2\int_0^\infty e^{-us}\,\frac{s^5\,\mathrm{d}s}{(s^2+1)^2},
\end{aligned}$$

which verifies the last sentence of the statement of Proposition 4.4; the second equality in the above display was obtained by taking the integral $\int_0^\infty e^{-us}w_2(s)\,u\,\mathrm{d}s$ by parts. Moreover, it follows that $g_1(u)$ is decreasing in $u > 0$. At that, $g_1(\infty-) = 0$, since $\mathcal{E}(\infty-) = 0$. So, on $(0,\infty)$ one has the following: $g_1 > 0$ and hence $g' > 0$, and therefore g is increasing. Since g is even and obviously continuous, it follows that indeed $g(u)$ increases in $|u|$. Clearly, $g(0) = |I(0)|^2 = 0$. It remains only to show that $g(\infty-) = 1$. Toward that end, integrate by parts to obtain the recursive relation $\mathcal{E}_j(u) = -ie^{-iu}u^{-j} + ij\mathcal{E}_{j+1}(u)$ for all $j > 0$. In particular, it follows that $\mathcal{E}_3(u) = -ie^{-iu}u^{-3} + 3i\mathcal{E}_4(u)$ and $\mathcal{E}_4(u) \precsim u^{-4} + |\mathcal{E}_5(u)| \leqslant u^{-4} + \int_u^\infty \frac{\mathrm{d}z}{z^5} \precsim u^{-4}$. Thus, $\mathcal{E}(u) = \mathcal{E}_3(u) = -ie^{-iu}u^{-3} + O(u^{-4}) = -iu^{-3}\big(e^{-iu} + o(1)\big)$ and $g(u) = u^6\,|\mathcal{E}(u)|^2 \to 1$ as $u \to \infty$. □

One will also could use a better upper bound on $|f(t)|$ for a given real value of t, where f is the c.f. of a r.v. X, say with $\mathsf{E}\,X = 0$, $\mathsf{E}\,X^2 = 1$, and a given value of $\rho := \mathsf{E}\,|X|^3$. Since

$$|f(t)| = \sqrt{\mathsf{E}^2\cos tX + \mathsf{E}^2\sin tX} = \sup_{\theta\in[0,2\pi]}\,(\cos\theta\ \mathsf{E}\cos tX + \sin\theta\ \mathsf{E}\sin tX) = \sup_{\theta\in[0,2\pi]}\,\mathsf{E}\cos(tX-\theta), \tag{4.22}$$

the best upper bound on $|f(t)|$ under the given conditions is

$$\begin{aligned}\mathcal{S}(t,\rho) &:= \sup\{|f(t)|\colon\ \mathsf{E}\,X=0, \mathsf{E}\,X^2=1, \mathsf{E}\,|X|^3=\rho\} \qquad (4.23)\\ &= \sup\{\mathsf{E}\cos(tX-\theta)\colon\ \mathsf{E}\,X=0, \mathsf{E}\,X^2=1, \mathsf{E}\,|X|^3=\rho, \operatorname{card}\operatorname{supp}X\leqslant 4, \theta\in[0,2\pi]\}\\ &= \sup\Big\{\sum_1^4 p_j\cos(tx_j-\theta)\colon\ \sum_1^4 p_j=1, \sum_1^4 p_jx_j=0, \sum_1^4 p_jx_j^2=1, \sum_1^4 p_j|x_j|^3=\rho,\\ &\qquad\qquad p_1,\dots,p_4\geqslant 0,\ \theta\in[0,2\pi]\Big\};\end{aligned}$$

card supp denotes the cardinality of the support of (the distribution of) X; for the second equality here, one can use the known results by Hoeffding [11] or Karr [12] or, somewhat more conveniently, Winkler [46] or Pinelis [31, Propositions 5 and 6(v)]. Thus the optimization problem reduces to one in 9 variables: $p_1,\dots,p_4,x_1,\dots,x_4,\theta$; in fact, one can easily solve the linear (or, more precisely, affine) restrictions $\sum_1^4 p_j=1, \sum_1^4 p_jx_j=0, \sum_1^4 p_jx_j^2=1, \sum_1^4 p_j|x_j|^3=\rho$ for $p_1,\dots,p_4$, and then only 5 variables will remain: $x_1,\dots,x_4,\theta$, with the additional restrictions on $x_1,\dots,x_4$ to provide for the conditions $p_1,\dots,p_4\geqslant 0$. For any given pair of values of (t,ρ), it will not be overly hard to find a close upper bound on the supremum $\mathcal{S}(t,\rho)$. A difficulty here is that one has to deal with two parameters, t and ρ, and obtain a close majorant of $\mathcal{S}(t,\rho)$ with, at least, discoverable and tractable patterns of monotonicity/convexity in t and ρ, if not with a more or less explicit expression. Apparently the main difficulty in dealing with $\mathcal{S}(t,\rho)$ will be that the target function $\cos(t\#-\theta)$ oscillates, whereas the function $(\#-w)_+^3$ in [34, Lemma 3.4] is monotonic.

Similar methods can be used to find a good upper bound on $|f(t)-g(t)|$, where $g=e^{-\#^2/2}$, the c.f. of the standard normal r.v. Z; in particular, one can start here by writing

$$|f(t)-g(t)| = \sup_{\theta\in[0,2\pi]}\big(\mathsf{E}\cos(tX-\theta)-\mathsf{E}\cos(tZ-\theta)\big) = \sup_{\theta\in[0,2\pi]}\big(\mathsf{E}\cos(tX-\theta)-g(t)\cos\theta\big)$$

in place of (4.22). At this point, one also has an option to use Stein's method to bound $\mathsf{E}\cos(tX-\theta)-\mathsf{E}\cos(tZ-\theta)$.

5. Constructions of the smoothing filter M

The following proposition was somewhat implicit in the paper [3] by Bohman.

Proposition 5.1. *Let p be any symmetric probability density function (p.d.f.) such that the function $\#p(\#)$ is integrable on $\mathbb{R}$. Take any real*

$$\kappa\geqslant\kappa_* := \frac{1}{\int_{\mathbb{R}}|x|p(x)\,\mathrm{d}x}. \qquad (5.1)$$

Let $\hat p$ stand, as usual, for the Fourier transform of p $\big($so that $\hat p(\#)=\int_{\mathbb{R}}e^{ix\#}p(x)\,\mathrm{d}x\big)$, and let then $\hat p'$ denote the derivative of $\hat p$ $\big($which exists, since $\int_{\mathbb{R}}|x|p(x)\,\mathrm{d}x<\infty\big)$. Then the function

$$M := \hat p + i\kappa\hat p' \qquad (5.2)$$

is such that inequalities (2.2) hold for all r.v. X, all real $T>0$, and all real x.

Because of the symmetry of p, in the conditions of Proposition 5.1 the function $\hat p=\operatorname{Re}M$ is even and $\kappa\hat p'=\operatorname{Im}M$ is odd, so that conditions (2.5) hold. In order to satisfy the conditions (2.1) and $M\in C^3$ as well, one may choose the symmetric p.d.f. $p_{0,2}$ defined by the formula

$$p_{0,2}(x) := \frac{32\pi^3}{3}\,\frac{1-\cos x}{x^2\left(x^2-4\pi^2\right)^2} \qquad (5.3)$$

for real $x \notin \{-2\pi, 0, 2\pi\}$ and then let M be as in (5.2) with $p = p_{0,2}$:

$$M = M_{0,2} := \widehat{p_{0,2}} + i\kappa \widehat{p_{0,2}}\,', \tag{5.4}$$

with any

$$\kappa \geqslant \kappa_{0,2} := \frac{1}{\int_{\mathbb{R}} |x| p_{0,2}(x)\, \mathrm{d}x} = 0.3418\dots. \tag{5.5}$$

Then $M_{0,2} \in C^3$, since $M''' = \hat{p}''' + i\kappa\hat{p}''''$ and

$$\int_{\mathbb{R}} x^4\, p_{0,2}(x)\, \mathrm{d}x < \infty. \tag{5.6}$$

Moreover, it is clear that $p_{0,2}$ is the restriction to $\mathbb{R}$ of an entire analytic function of exponential type 1; so, by the Paley–Wiener theory (see e.g. [7, Section 43]), the condition (2.1) holds as well, with $M_{0,2}$ in place of M. In fact,

$$\operatorname{Re} M_{0,2}(t) = \widehat{p_{0,2}}(t) = \left(\frac{2 + \cos 2\pi t}{3}\,(1 - |t|) + \frac{\sin 2\pi|t|}{2\pi}\right) \mathrm{I}\{|t| < 1\} \quad \text{and} \tag{5.7}$$

$$\operatorname{Im} M_{0,2}(t) = \kappa \widehat{p_{0,2}}\,'(t) = -\kappa\, \frac{\operatorname{sign} t}{3} \left[2\pi(1 - |t|)\sin 2\pi|t| + 4\sin^2(\pi t)\right] \mathrm{I}\{|t| < 1\} \tag{5.8}$$

for all real t. Graphs of $p_{0,2}$, $\operatorname{Re} M_{0,2}$, and $\operatorname{Im} M_{0,2}$ with $\kappa = \kappa_{0,2}$ are shown in Fig. 2.



Fig 2: Graphs of $p_{0,2}$, $\operatorname{Re} M_{0,2}$, and $\operatorname{Im} M_{0,2}$ with $\kappa = \kappa_{0,2}$.

More generally, in order that a function M as in (5.2) satisfy the conditions (2.1) and $M \in C^3$, it is enough that $\hat{p}$ be smooth enough and such that (2.1) holds with $\hat{p}$ in place of M. Therefore, the following well-known characterization is useful.

Proposition 5.2. *(See e.g. [14, Theorem 4.2.4].) A function $f\colon \mathbb{R} \to \mathbb{C}$ is the c.f. of an absolutely continuous distribution on $\mathbb{R}$ if and only if $f(0) = 1$ and $f = g * \overline{g}^-$ for some (possibly complex-valued) function $g \in L^2(\mathbb{R})$. Here and in the sequel, as usual, the symbol $*$ stands for the convolution, the bar denotes the complex conjugation, $g^-(\#) := g(-\#)$, and $\overline{g}^- := (\overline{g})^- = \overline{g^-}$.*

Indeed, take any smooth enough nonzero function $g\colon \mathbb{R} \to \mathbb{C}$ such that $g(t) = 0$ for all real $t \notin [a, b]$, where a and b are any real numbers such that $a < b$. Then, by Proposition 5.2, the function $f := g * \overline{g}^- / \|g\|_2^2$ is the c.f. of an absolutely continuous distribution on $\mathbb{R}$, f is smooth enough, and $f(t) = 0$ for all real $t \notin [-T, T]$, where $T := b - a$. At that, if g is real-valued, then f is even. To spell-out the “smooth enough” condition and conclusion here, one can easily check that, if $g \in C^j$ and $h \in C^k$ for some j and k in $\mathbb{N}$ and (say) $|g(t)| + |h(t)| = 0$ for some real $T > 0$ and all real $t \notin [0, T]$, then $g * h \in C^{j+k}$, with $(g * h)^{(j+k)} = g^{(j)} * h^{(k)}$.

One can use Proposition 5.1 and Proposition 5.2 to optimize properties of the filter M – say by taking g to be an arbitrary nonzero real-valued spline of a high enough order and/or with a large enough subintervals of the interval $[0,T]$, extending g to $\mathbb{R}$ by letting $g(t) := 0$ for all real $t \notin [0,T]$, letting then $f := g * \overline{g}^-/\|g\|_2^2$, defining M as in (5.2), and finally (quasi-)optimizing with respect to the parameters of the spline.

While the construction described in Proposition 5.1 is comparatively simple, it appears somewhat too rigid and wasteful. Indeed, in order that the imaginary part $M_2 = \kappa\hat{p}'$ of the function M in (5.2) be thrice differentiable $\big($as needed or almost needed in the *quick proof* beginning on page 10$\big)$, the real part $M_1 = \hat{p}$ of M must be four times differentiable; equivalently $\big($cf. (5.6)$\big)$, the density p must have light enough tails so that $\int_{\mathbb{R}} x^4\, p(x)\,\mathrm{d}x < \infty$. Together with the filtering condition (2.1), the condition of extra smoothness of $M_1 = \hat{p}$/extra lightness of the tails of p may result in a smoothing filter M which is not as good as it can be, thus compromising the quality of the approximation by the upper and lower bounds in (4.9).

A more flexible and potentially better construction of the smoothing filter M can be given as follows. As in Proposition 5.1, let us start with an arbitrary symmetric p.d.f. p, whose Fourier transform $\hat{p}$ is intended to be $M_1 = \operatorname{Re} M$. Accordingly, let us assume right away that

$$\hat{p}(t) = 0 \quad \text{if} \quad |t| > 1; \tag{5.9}$$

cf. (2.1). Note that the smoothing filter as in (5.2) is the Fourier transform of the function

$$x \mapsto p(x)(1-\kappa x), \tag{5.10}$$

which differs relatively much from the "original" p.d.f. $x \mapsto p(x)$ when $|x|$ is large. To address this concern, let us replace the "large" factor x in (5.10) by a "tempered" and, essentially, more general factor $G(x)$ such that $G\colon \mathbb{R} \to \mathbb{R}$ is a strictly increasing odd function of bounded variation, whose Fourier–Stieltjes transform $\widehat{\mathrm{d}G}(\#) = \int_{\mathbb{R}} e^{ix\#}\,\mathrm{d}G(x)$ satisfies the condition

$$\widehat{\mathrm{d}G}(t) = 0 \quad \text{if} \quad |t| > \gamma, \tag{5.11}$$

for some real $\gamma > 0$. The no-high-frequency-component condition (5.11) implies, by the mentioned Paley–Wiener theory, that the function G is the restriction to $\mathbb{R}$ of an entire analytic function of exponential type γ and, in particular, is infinitely many times differentiable. Without loss of generality, assume that the function $\frac{1}{2} + G$ is a d.f.

As mentioned above, instead of the "harsh" tilting (5.10) of the p.d.f. p, we consider the "tempered" tilting:

$$x \mapsto \tilde{p}(x) := p(x)\big(1 - \kappa G(x)\big), \tag{5.12}$$

for any real

$$\kappa \geqslant \kappa_* := \frac{1}{2\int_0^\infty p(x)G(x)\,\mathrm{d}x}. \tag{5.13}$$

Note that $0 \leqslant G(x) < \frac{1}{2} = G(\infty-)$ for all real $x > 0$; also, as discussed previously, the condition (5.9) implies that p is the restriction to $\mathbb{R}$ of an entire analytic function, and so, $p > 0$ almost everywhere on $\mathbb{R}$. Therefore and by the symmetry of p, one has $0 < \int_0^\infty p(x)G(x)\,\mathrm{d}x < \frac{1}{2}\int_0^\infty p(x)\,\mathrm{d}x = \frac{1}{4}$ and hence $\kappa_* > 2$ and $\kappa > 2$. It follows that there exists a unique root $x_\kappa \in (0,\infty)$ of the equation

$$1 - \kappa G(x_\kappa) = 0. \tag{5.14}$$

Hence, $\tilde{p} \geqslant 0$ on the interval $(-\infty, x_\kappa]$ and $\tilde{p} \leqslant 0$ on $[x_\kappa, \infty)$, so that the function

$$\tilde{F}(\#) := \int_{-\infty}^{\#} \tilde{p}(y)\,\mathrm{d}y \tag{5.15}$$

is nondecreasing on $(-\infty, x_\kappa]$ and nonincreasing on $[x_\kappa, \infty)$. At that, $\tilde{F}(\infty-) = 1$, since p is an even p.d.f. and the bounded function G is odd; also, clearly $\tilde{F}(-\infty+) = 0$. Moreover, $\tilde{F}(0) = \frac{1}{2} - \kappa \int_{-\infty}^{0} p(x)G(x)\,\mathrm{d}x = \frac{1}{2} + \kappa \int_{0}^{\infty} p(x)G(x)\,\mathrm{d}x = \frac{1}{2} + \frac{\kappa}{2\kappa_*} \geqslant 1$. It follows that

$$\mathrm{I}\{y \geqslant 0\} \leqslant \tilde{F}(y) \quad \text{for all real } y. \tag{5.16}$$

Let now X be any r.v. and let f by its c.f.:

$$f(\#) := \mathsf{E}\, e^{iX\#}. \tag{5.17}$$

Then $\mathsf{P}(X \leqslant x) = \mathsf{E}\,\mathrm{I}\{x - X \geqslant 0\} \leqslant \mathsf{E}\,\tilde{F}(x - X)$, by (5.16); that is,

$$\mathsf{P}(X \leqslant x) \leqslant \int_{\mathbb{R}} \tilde{F}(x - y)\,\mathsf{P}(X \in \mathrm{d}y), \tag{5.18}$$

for all real x. Define now M as the Fourier transform of $\tilde{p}$, so that

$$M = \hat{\tilde{p}}, \quad M_1 = \operatorname{Re} M = \hat{p}, \quad \text{and} \quad M_2 = \operatorname{Im} M = i\kappa\,\widehat{pG}, \tag{5.19}$$

by (5.12). Note that the Fourier–Stieltjes transform of the function $\int_{\mathbb{R}} \tilde{F}(\# - y)\,\mathsf{P}(X \in \mathrm{d}y)$ is the Fourier transform of $\int_{\mathbb{R}} \tilde{p}(\# - y)\,\mathsf{P}(X \in \mathrm{d}y)$, which in turn is $\hat{\tilde{p}} f = Mf$. Then, in view of Proposition 4.2, (5.18) means that the last inequality in (2.2) holds for $T = 1$; that it holds for any real $T > 0$ now follows by simple re-scaling, since (5.16) obviously implies $\mathrm{I}\{y \geqslant 0\} \leqslant \tilde{F}(Ty)$ for all real y and all $T > 0$. Similarly or using the reflection $x \mapsto -x$, one can see that the first inequality in (2.2) holds as well.

To compute M_2 in (5.19), we need to express $\widehat{pG}$ in terms $\hat{p}$ and $\widehat{\mathrm{d}G}$. To simplify the derivation, assume the condition $\hat{p} \in C^1$, as well as the previously stated conditions (5.9) and (5.11); these conditions will hold in the applications anyway. Then one can see that for all real u

$$\widehat{pG}(u) = \frac{i}{2\pi}\,\text{p.v.}\int_{-\infty}^{\infty} \hat{p}(u - s)\,\widehat{\mathrm{d}G}(s)\,\frac{\mathrm{d}s}{s} = \frac{i}{2\pi}\int_{\mathbb{R}} \frac{\hat{p}(u - s) - \hat{p}(u)}{s}\,\widehat{\mathrm{d}G}(s)\,\mathrm{d}s; \tag{5.20}$$

the latter equality here holds because the function G was assumed odd, and hence the function $\widehat{\mathrm{d}G}$ is even; the latter integral in (5.20) may be understood in the Lebesgue sense, in view of the conditions $\hat{p} \in C^1$ and (5.11). It follows from (5.20), (5.9), and (5.11) that

$$\widehat{pG}(t) = 0 \quad \text{if} \quad |t| > 1 + \gamma. \tag{5.21}$$

To verify the first equality in (5.20), one can write

$$\begin{aligned} p(x)G(x) &= \Big(\frac{1}{2\pi}\int_{\mathbb{R}} e^{-itx}\hat{p}(t)\,\mathrm{d}t\Big)\Big(\frac{i}{2\pi}\,\text{p.v.}\int_{-\infty}^{\infty} e^{-isx}\,\widehat{\mathrm{d}G}(s)\,\frac{\mathrm{d}s}{s}\Big) \\ &= \frac{1}{2\pi}\int_{\mathbb{R}} e^{-iux}\,\mathrm{d}u\,\frac{i}{2\pi}\,\text{p.v.}\int_{-\infty}^{\infty} \hat{p}(u - s)\,\widehat{\mathrm{d}G}(s)\,\frac{\mathrm{d}s}{s}; \end{aligned}$$

the second equality here is justified because of the second equality in (5.20) and the inequality $\big|\int_{\mathbb{R}} \frac{\hat{p}(u-s)-\hat{p}(u)}{s}\,\widehat{\mathrm{d}G}(s)\,\mathrm{d}s\big| \leqslant 2\gamma\,\mathrm{I}\{|u| \leqslant 1 + \gamma\}\max_{|t|\leqslant 1}|\hat{p}'(t)|$ for all real u.

Note that the first integral in (5.20) is a convolution. One can also integrate by parts to represent $\widehat{pG}$ as a convolution-smoothing of the derivative $\hat{p}'$:

$$\widehat{pG} = \tfrac{i}{2\pi}\,\hat{p}' * \widetilde{\widehat{\mathrm{d}G}}, \tag{5.22}$$

where

$$\widetilde{\widehat{\mathrm{d}G}}(t) := \text{p.v.}\int_{-\infty}^{t} \widehat{\mathrm{d}G}(s)\frac{\mathrm{d}s}{s} := \lim_{\varepsilon\downarrow 0}\int_{(-\infty,t)\setminus(-\varepsilon,\varepsilon)} \widehat{\mathrm{d}G}(s)\frac{\mathrm{d}s}{s} = \int_{-\gamma}^{-|t|} \widehat{\mathrm{d}G}(s)\frac{\mathrm{d}s}{s}\ \mathrm{I}\{|t|<\gamma\} \tag{5.23}$$

for all real t. This follows because for all real u

$$\begin{aligned}\int_{\mathbb{R}} \frac{\hat{p}(u-s)-\hat{p}(u)}{s}\,\widehat{\mathrm{d}G}(s)\,\mathrm{d}s &= \int_{\mathbb{R}} \widehat{\mathrm{d}G}(s)\frac{\mathrm{d}s}{s}\Big(\int_{u}^{u-s} \hat{p}'(v)\,\mathrm{d}v\ \mathrm{I}\{s<0\} - \int_{u-s}^{u} \hat{p}'(v)\,\mathrm{d}v\ \mathrm{I}\{s>0\}\Big) \\ &= \int_{u}^{\infty} \hat{p}'(v)\,\mathrm{d}v \int_{-\infty}^{u-v} \widehat{\mathrm{d}G}(s)\frac{\mathrm{d}s}{s} - \int_{-\infty}^{u} \hat{p}'(v)\,\mathrm{d}v \int_{u-v}^{\infty} \widehat{\mathrm{d}G}(s)\frac{\mathrm{d}s}{s} \\ &= \int_{\mathbb{R}} \hat{p}'(v)\,\mathrm{d}v\,\widetilde{\widehat{\mathrm{d}G}}(u-v),\end{aligned}$$

since the function $\widehat{\mathrm{d}G}$ is even. The latter condition or (5.23) also shows that the function $\widetilde{\widehat{\mathrm{d}G}}$ is even. Moreover, since $\widehat{\mathrm{d}G}(s)\to 1$ as $s\to 0$, (5.23) yields

$$\widetilde{\widehat{\mathrm{d}G}}(t) \sim \ln|t| \tag{5.24}$$

as $t\to 0$; thus, the function $\widetilde{\widehat{\mathrm{d}G}}$ is mildly singular in a neighborhood of 0. For instance, $\widetilde{\widehat{\mathrm{d}G}}(t) \equiv (1-|t|+\ln|t|)(\mathrm{I}\{|t|<1\}$ if $\widehat{\mathrm{d}G}(t) \equiv (1-|t|)(\mathrm{I}\{|t|<1\}$.

Note also that in the case $\big($prevented by the condition (5.11)$\big)$ when $\widehat{\mathrm{d}G}=1$ on $\mathbb{R}$, the function $-2i\,\widehat{pG}$ would be the Hilbert transform of the function $\hat{p}$; see e.g. [42, Chapter V].

It follows from (5.19) and (5.21) that

$$M_2(t)=0 \quad \text{if} \quad |t|>1+\gamma. \tag{5.25}$$

This condition on M_2 is obviously weaker than the condition

$$M_2(t)=0 \quad \text{if} \quad |t|>1, \tag{5.26}$$

following from (2.1) and (2.5). However, by (5.19) and (5.9), one still has $M_1(t)=0$ if $|t|>1$, whereas the condition (5.26) was used in the *quick proof* beginning on page 10 only to bound two terms, $\mathfrak{G}_{2\alpha}(f_{31})$ and $\mathfrak{G}_{2\alpha}(f_{32})$. Therefore, one may expect the adverse impact of the weakening of the condition (5.26) to (5.25) to be rather limited and likely more than compensated for by the advantages provided by the more flexible construction of the smoothing filter M, with the tempered tilting of M_1. Moreover, the latter construction is, essentially, more general, Indeed, for instance, one may always include G into the scale family $(G_\alpha)_{\alpha>0} := \big(G(\frac{\#}{\alpha})\big)_{\alpha>0}$, and then the tempered tilting (5.12) will be close to the harsh tilting (5.10) for large $\alpha>0$ provided that $G'(0)\neq 0$. Indeed, $G_\alpha(x)\sim\frac{G'(0)}{\alpha}x$ for each real $x\neq 0$ as $\alpha\to\infty$; of course, at that the value of $\kappa=\kappa_\alpha$ in (5.12) with $G=G_\alpha$ will be quite different from that in (5.10); in fact, the value of κ_* in (5.13) with $G=G_\alpha$ will then be asymptotically equivalent to the value of κ_* in (5.1) times $\frac{\alpha}{G'(0)}$, provided that $\int_{\mathbb{R}}|x|p(x)\,\mathrm{d}x<\infty$. At that, the value of γ in (5.11) for G will be replaced by the corresponding value $\gamma_\alpha := \frac{\gamma}{\alpha}$ for G_α, so that $\gamma_\alpha\to 0$ as $\alpha\to\infty$.

References

[1] V. Bentkus. On the asymptotical behavior of the constant in the Berry-Esseen inequality. *J. Theoret. Probab.*, 7(2):211–224, 1994.

[2] A. Bikelis. Estimates of the remainder term in the central limit theorem. *Litovsk. Mat. Sb.*, 6:323–346, 1966.

[3] H. Bohman. To compute the distribution function when the characteristic function is known. *Skand. Aktuarietidskr.*, 1963:41–46 (1964), 1964.

[4] L. H. Y. Chen and Q.-M. Shao. Stein's method for normal approximation. In *An introduction to Stein's method*, volume 4 of *Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap.*, pages 1–59. Singapore Univ. Press, Singapore, 2005.

[5] G. P. Chistyakov. On a problem of A. N. Kolmogorov. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 184(Issled. po Mat. Statist. 9):289–319, 326–327, 1990.

[6] T. Dinev and L. Mattner. The asymptotic Berry-Esseen constant for intervals. `http://arxiv.org/abs/1111.7146`, 2011.

[7] W. F. Donoghue, Jr. *Distributions and Fourier transforms.* Academic Press, New York, 1969. Pure and Applied Mathematics, Volume 32.

[8] C. G. Esseen. A moment inequality with an application to the central limit theorem. *Skand. Aktuarietidskr.*, 39:160–170 (1957), 1956.

[9] S. V. Gavrilenko. Refinement of nonuniform estimates of the rate of convergence of the distributions of poisson random sums to the normal law (in Russian). *Informatika i ee Primeneniya (Informatics and Applications)*, 5:12–24, 2011.

[10] J. Gurland. Inversion formulae for the distribution of ratios. *Ann. Math. Statistics*, 19:228–237, 1948.

[11] W. Hoeffding. The extrema of the expected value of a function of independent random variables. *Ann. Math. Statist.*, 26:268–275, 1955.

[12] A. F. Karr. Extreme points of certain sets of probability measures, with applications. *Math. Oper. Res.*, 8(1):74–85, 1983.

[13] V. Y. Korolev and I. G. Shevtsova. An improvement of the Berry–Esseen inequality with applications to Poisson and mixed Poisson random sums. *Scandinavian Actuarial Journal*, 2012(2):81–105, 2012.

[14] E. Lukacs. *Characteristic functions.* Hafner Publishing Co., New York, 1970. Second edition, revised and enlarged.

[15] R. Michel. On the constant in the nonuniform version of the Berry-Esseen theorem. *Z. Wahrsch. Verw. Gebiete*, 55(1):109–117, 1981.

[16] S. A. Mirakhmedov. On the absolute constant in a nonuniform estimate of the rate of convergence in the central limit theorem. *Izv. Akad. Nauk UzSSR Ser. Fiz.-Mat. Nauk*, (4):26–30, 78, 1984.

[17] A. V. Nagaev. Integral limit theorems with regard to large deviations when Cramér's condition is not satisfied. I. *Theor. Probability Appl.*, 14:51–64, 1969.

[18] A. V. Nagaev. Integral limit theorems with regard to large deviations when Cramér's condition is not satisfied. II. *Theor. Probability Appl.*, 14:193–208, 1969.

[19] S. V. Nagaev. Some limit theorems for large deviations. *Teor. Verojatnost. i Primenen*, 10:231–254, 1965.

[20] Y. S. Nefedova and I. G. Shevtsova. On the accuracy of normal approximation for the distributions of poisson random sums (in Russian). *Informatika i ee Primeneniya (Informatics and Applications)*, 5(1):39–45, 2011.

[21] Y. S. Nefedova and I. G. Shevtsova. Nonuniform estimates of convergence rate in the central limit theorem (in Russian). *Teor. Veroyatnost. i Primenen.*, 57(1):62–97, 2012.

[22] L. Paditts and S. A. Mirakhmedov. Letter to the editors: "On the absolute constant in a nonuniform estimate of the rate of convergence in the central limit theorem" [Izv. Akad. Nauk UzSSR Ser. Fiz.-Mat. Nauk **1984**, no. 4, 26–30; MR MR0780094 (86i:60071)]. *Izv. Akad. Nauk*

UzSSR Ser. Fiz.-Mat. Nauk, (3):80, 1986.

[23] L. Paditz. *Über die Annäherung der Verteilungsfunktionen von Summen unabhängiger Zufallsgröen gegen unbegrenzt teilbare Verteilungsfunktionen unter besonderer beachtung der Verteilungsfunktion der standardisierten Normalverteilung. Dissertation A.* PhD thesis, Technische Universität Dresden, 1977.

[24] L. Paditz. Abschätzungen der Konvergenzgeschwindigkeit zur Normalverteilung unter Voraussetzung einseitiger Momente. *Math. Nachr.*, 82:131–156, 1978.

[25] L. Paditz. On the analytical structure of the constant in the nonuniform version of the Esseen inequality. *Statistics*, 20(3):453–464, 1989.

[26] V. V. Petrov. *Sums of independent random variables.* Springer-Verlag, New York, 1975. Translated from the Russian by A. A. Brown, Ergebnisse der Mathematik und ihrer Grenzgebiete, Band 82.

[27] R. R. Phelps. *Lectures on Choquet's theorem.* D. Van Nostrand Co., Inc., Princeton, N.J.-Toronto, Ont.-London, 1966.

[28] I. Pinelis. Improved nonuniform Berry–Esseen-type bounds, preprint, `http://arxiv.org/find/all/1/au:+pinelis/0/1/0/all/0/1`.

[29] I. Pinelis. Exact bounds on the truncated-tilted mean, with applications (preprint). `http://arxiv.org/abs/1103.3683`, 2011.

[30] I. Pinelis. Exact lower bounds on the exponential moments of Winsorized and truncated random variables. *J. App. Probab.*, 48:547–560, 2011.

[31] I. Pinelis. On the extreme points of moments sets. `http://arxiv.org/abs/1204.0249`, 2011.

[32] I. Pinelis. Optimal re-centering bounds, with applications to Rosenthal-type concentration of measure inequalities. `http://arxiv.org/abs/1111.2622`, to appear in Proceedings of the conference High Dimensional Probability VI, Banff, October 9–14, 2011, 2011.

[33] I. Pinelis. An exact bound on the truncated-tilted mean for symmetric distributions. `http://arxiv.org/abs/1205.5234`, 2012.

[34] I. Pinelis. On the Bennett-Hoeffding inequality, to appear in *Annales de l'Institut Henri Poincaré.* 2012.

[35] I. F. Pinelis. A problem on large deviations in a space of trajectories. *Theory Probab. Appl.*, 26(1):69–84, 1981.

[36] I. F. Pinelis. Asymptotic equivalence of the probabilities of large deviations for sums and maximum of independent random variables. In *Limit theorems of probability theory*, volume 5 of *Trudy Inst. Mat.*, pages 144–173, 176. "Nauka" Sibirsk. Otdel., Novosibirsk, 1985.

[37] I. F. Pinelis and S. A. Utev. Sharp exponential estimates for sums of independent random variables. *Theory Probab. Appl.*, 34(2):340–346, 1989.

[38] H. Prawitz. Limits for a distribution, if the characteristic function is given in a finite domain. *Skand. Aktuarietidskr.*, pages 138–154 (1973), 1972.

[39] I. Shevtsova. On the absolute constants in the Berry-Esseen type inequalities for identically distributed summands. `http://arxiv.org/abs/1111.6554`, 2011.

[40] I. G. Shevtsova. Refinement of estimates for the rate of convergence in Lyapunov's theorem. *Dokl. Akad. Nauk*, 435(1):26–28, 2010.

[41] I. S. Shiganov. Refinement of the upper bound of a constant in the remainder term of the central limit theorem. In *Stability problems for stochastic models (Moscow, 1982)*, pages 109–115. Vsesoyuz. Nauchno-Issled. Inst. Sistem. Issled., Moscow, 1982.

[42] E. C. Titchmarsh. *Introduction to the theory of Fourier integrals.* Chelsea Publishing Co., New York, third edition, 1948.

[43] W. Tysiak. *Gleichmäige und nicht-gleichmige Berry-Esseen-Abschätzungen. Dissertation.* PhD thesis, Universität-Gesamthochschule Wuppertal, 1983.

[44] I. Tyurin. New estimates of the convergence rate in the Lyapunov theorem (preprint, arXiv:0912.0726v1 [math.PR]).

[45] J. D. Vaaler. Some extremal functions in Fourier analysis. *Bull. Amer. Math. Soc. (N.S.)*, 12(2):183–216, 1985.

[46] G. Winkler. Extreme points of moment sets. *Math. Oper. Res.*, 13(4):581–587, 1988.